

Iron

PLANT ACTIVE CHELATE FORMULATION



THE PRODUCT

 Amino Boss Iron is an iron nutrition deficiency corrector for all crops. It can be used in a preventive program or at the onset of symptoms.

DIRECTIONS FOR USE

- Product should be used during ideal growing conditions to ensure effective uptake. Avoid extreme weather conditions such as heat and frost events.
 Spraying in the early morning or late afternoon can assist in minimising risk of rapid drying on the leaf due to evaporation. Where local experience of successful use is not available, we recommend a small scale jar test and test application prior to wide spread use. If any issues are observed do not continue with the application.
- It is recommended to add product to the tank mix after all other products have been mixed well in the tank, including most of the water.
- Do not apply by foliar application to stone fruit.
- Do not apply by foliar application to flowers or fruit of any crop.

COMPATIBILITY

Compatible with most common fertilizers and agrochemicals used. It is recommended to perform a jar test before mixing. Do not use with amine agricultural chemicals. Do not apply **Amino Boss Iron** with copper fungicides by foliar application on all crops.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre, Phone Australia 131 126.

SAFETY DATA SHEET

For further information, refer to the Safety Data Sheet (SDS) which is available from the supplier or from our web site, **www.sipcam.com.au**

Highly available plant active iron formulation Total iron (Fe) content of 5% w/v as Amino Acid Chelate

A unique formulation of Iron amino acid chelate solution containing the complete range of amino acids to assist in chelation and plant performance. Recommended for foliar applications where a 100% chelated product with excellent plant uptake and good compatibility is required.

CROPS	RATE	COMMENTS
BROADACRE CROPS	Foliar 1 – 2 L/ha	Apply Cereals 3-5 leaf stage. Pulses/Lucerne 10-14 days before flower Canola 4-9 true leaves
TREES (excluding stonefruit), NUTS, BERRIES AND VINES	Foliar 1 – 3 L/ha Fertigation 2 – 5 L/ha	Apply in a water volume of 400L to 1000L per ha. Apply at regular intervals of 7-14 days from petal fall up until harvest.
VEGETABLES Cucurbits Fruiting Vegetables Potatoes Carrots, Onions	Foliar & Fertigation 500mL – 2 L/ha	Apply from onset of active growth

CAUTION: Do not apply as a foliar application in conjunction with copper and EC fungicides. As **Amino Boss Iron** contains a strong chelating agent, it will increase the activity of copper and levels of resulting phytotoxicity may not be acceptable to growers. For most effective results apply early morning or late afternoon.

WARNING STATEMENTS

Do not swallow. The spray from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin.

Batch No: Date of Manufacture:

This product is NOT a Dangerous Good under the Australian Code for the transport of Dangerous Goods by road and rail

Specialist Advice In Emergency Only **1800 033 111** All Hours - Australia Wide



Sipcam Pacific Australia Pty Ltd Level 1, 191 Malop Street Geelong

1000 L