

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

BRIMSTONE[®]

HERBICIDE

ACTIVE CONSTITUENT: 240 g/L IMAZAPIC present as the ammonium salt



For the pre-emergent control of certain annual grass and broadleaf weeds in fallow situations, sugarcane and peanuts and early post-emergence control of certain annual grass and broadleaf weeds in peanuts and sugarcane as specified in the DIRECTIONS FOR USE table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT.

CONTENTS: 1L, 5L, 20L

APVMA Approval No: 69352/60574



SIPCAM PACIFIC AUSTRALIA PTY LIMITED
Level 1, 191 Malop Street Geelong VIC 3220

® Registered Trademark of Sipcam Pacific Australia Pty Ltd

BRIMSTONE HERBICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. When opening the container and preparing the spray, wear elbow length PVC gloves. Wash hands after use. After each days use, wash gloves.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. *Phone Australia 131126.*

SAFETY DATA SHEET

For further information, refer to the Safety Data Sheet (SDS) which is available from the supplier or from our web site, www.sipcam.com.au

NOTICE TO BUYER

This product must be used in accordance with the directions for use. Where this product is not a good of a kind ordinarily acquired for personal, domestic or household use or consumption under the Australian Consumer Law, Sipcam's liability for any direct, indirect or consequential loss which you incur, including, without limitation, loss or damage to crop, loss of equipment, property damage, personal injury or death is limited, at Sipcam's option, to the replacement of the product or the supply of an equivalent product. By opening this package, you agree to be bound by these terms. If you do not agree to these terms, please return the entire unopened package intact to Sipcam or the place in which you purchased this product for a full refund.

Batch No:

Date of Manufacture:

| |
|--|
| <p>Not subject to the provisions of the Australian Code for the Transport of Dangerous Goods by Road and Rail when packed in this container and transported in Australia.</p> |
|--|

| |
|--|
| <p>For specialist advice in emergency only, call 1800 033 111 all hours, Australia wide</p> |
|--|



READ SAFETY DIRECTIONS BEFORE OPENING OR USING

BRIMSTONE®

HERBICIDE

ACTIVE CONSTITUENT: 240 g/L IMAZAPIC present as the ammonium salt



For the pre-emergent control of certain annual grass and broadleaf weeds in fallow situations, sugarcane and peanuts and early post-emergence control of certain annual grass and broadleaf weeds in peanuts and sugarcane as specified in the DIRECTIONS FOR USE table.

IMPORTANT: READ THE THIS LEAFLET BEFORE USING THIS PRODUCT

CONTENTS: 1L, 5L, 20L

APVMA Approval No: 69352/60574



SIPCAM PACIFIC AUSTRALIA PTY LIMITED
Level 1, 191 Malop Street Geelong VIC 3220

® Registered Trademark of Sipcaml Pacific Australia Pty Ltd

DIRECTIONS FOR USE:**RESTRAINTS:**

DO NOT apply to soils saturated with water.

DO NOT apply by aircraft.

| SITUATION | WEEDS CONTROLLED | RATE per ha | CRITICAL COMMENTS |
|---|--|---------------|---|
| Fallow prior to planting winter crops as directed in the Follow Crops Table (Qld and Nthn NSW only) | Barnyard grass (<i>Echinochloa colona</i>), Liverseed grass (<i>Urochloa panicoides</i>), Stink grass (<i>Eragrostis ciliaris</i>), Blowaway grass (<i>Panicum decompositum</i>), Button grass (<i>Dactyloctenium radulans</i>), Pigweed (<i>Portulaca oleracea</i>), Yellow vine (<i>Tribulus terrestris</i>), Mintweed (<i>Salvia reflexa</i>), Boggabri weed (<i>Amaranthus mitchelli</i>), Dwarf amaranth (<i>Amaranthus macrocarpus</i>), Peachvine/cowvine (<i>Ipomoea lonchophylla</i>) | 150 to 200 mL | Treatments may be applied from the end of the previous crop harvest but not less than 3 months prior to planting of winter crop (refer to Follow Crops Table). Best results are obtained where treatments are applied to dry, weed-free soil prior to weed germination. Treatments can be applied to standing crop residues. Sufficient rainfall is required after application and prior to weed emergence to wet soil to a depth of 5 cm. Use the higher rates for increased residual control of weeds or anticipated high weed pressure. Emerged weeds must be controlled by prior cultivation, or by application of a knockdown herbicide. Where infestation of emerged weeds is light, tank mix treatment with glyphosate or paraquat. Cultivation following application of treatments may reduce pre-emergence weed control provided by Brimstone. Prolonged wet soil conditions following treatment will reduce the effective life of pre-emergence treatments. Weed escapes may require follow up application of knockdown herbicides. DO NOT apply more than 200 mL/ha per season on cropland. DO NOT apply in more than two successive seasons on cropland. DO NOT overspray headlands. |
| Sugarcane (Qld, NSW and WA only) | Awnless barnyard grass (<i>Echinochloa colona</i>), Barnyard grass (<i>E. crus-galli</i>), Blackberry nightshade (<i>Solanum nigrum</i>), Blue billygoat weed (<i>Ageratum houstonianum</i>), Common sida (<i>Sida rhombifolia</i>), Giant/Black pigweed (<i>Trianthema portulacastrum</i>), Glossy nightshade (<i>Solanum americanum</i>), Green amaranth (<i>Amaranthus viridis</i>), Green summer grass (<i>Brachiaria subquadriflora</i>), <i>Ipomoea</i> spp., Guinea grass (<i>Panicum maximum</i>), Liverseed grass (<i>Urochloa panicoides</i>), Milkweed (Mexican fire weed) (<i>Euphorbia heterophylla</i>), Pigweed (<i>Portulaca oleracea</i>), Summer grass (<i>Digitaria ciliaris</i>) | 300 to 400 mL | DO NOT apply more than once a year to the same crop. Pre-emergence application: PLANT CANE: Apply after planting and before emergence of the cane. On sandy soil, ensure thorough coverage of setts to a minimum depth of 5 cm. RATOON: Apply after harvest and before new cane growth appears. Use the higher rates of Brimstone for increased residual control of weeds or anticipated high weed pressure. Heavy trash layers may prevent the product from reaching the soil. Apply to weed-free soil prior to weed germination. Sufficient rainfall, or overhead irrigation, is required after application and prior to weed emergence to wet soil to a minimum depth of 5 cm. Emerged weeds must be controlled by prior cultivation, or application of an appropriate herbicide. Weed control can be improved under dry conditions or where incorporating moisture has been insufficient, by cultivating the treated soil above setts. Care needs to be taken to minimise exposure of untreated soil or setts. Pre-emergence weed control may be limited on Krasnozem or red brown Ferrosol soil types, where moisture is not maintained in the top 5 cm of soil. Refer to Soil Type in the General Instructions. Heavy rain and/or irrigation within 2 days following application may concentrate Brimstone in the furrow and cause temporary yellowing and stunting of cane leaves. Prolonged wet soil and/or cool conditions can increase crop damage, especially in sandy soil. Normal growth will return when conditions become warm and drier. Weed escapes may require follow up application of an appropriate herbicide. Contact Sipcam for advice on variety susceptibility before using this product (refer to the CROP SAFETY section of the label). |

| SITUATION | WEEDS CONTROLLED | RATE per ha | CRITICAL COMMENTS |
|----------------------------------|---|---|--|
| Sugarcane (Qld, NSW and WA only) | As above plus Crowsfoot grass (<i>Eleusine indica</i>) | 400 mL plus 3L pendimethalin 330g/L | Incorporate into the soil within 7 days of application, either by overhead irrigation, rainfall or mechanical incorporation, to a minimum depth of 5 cm. Read all directions for use in sugarcane as well as the Restraints and Follow Crops restrictions on the label for pendimethalin. |
| | As above plus control of those weeds expected with these rates of 250 g/L paraquat products alone. | 300 to 400 mL plus 1.2 to 1.6 L of 250 g/L paraquat product (e.g. Inferno) | Post-emergence application (in emerged cane): ALWAYS tank mix Brimstone with registered and appropriate rates of paraquat for the emerged weeds. Use the higher rates of Brimstone for increased residual control of weeds or anticipated high weed pressure. Use the higher rates of paraquat for dense, more mature weeds. Apply as a carefully directed inter-row spray to weeds and at the base of the plant, minimising contact with sugarcane foliage. Contact with sugarcane foliage may cause temporary crop damage observed as inter-veinal yellowing and reduction in crop biomass or crop height. Refer to Inferno (or equivalent) label re spray boom types and precautions. Reduce the incidence of "fines" by using the correct nozzles and operating pressures. Replace worn or old nozzles and calibrate equipment on a regular basis. DO NOT cultivate within 1 hour of application. Heavy rain and/or irrigation within 2 days following application may concentrate Brimstone in the furrow and cause temporary yellowing and stunting of cane leaves. DO NOT spray plants covered with heavy dew. Prolonged wet soil and/or cool conditions can increase crop damage, especially in sandy soil. Normal growth will return when conditions become warm and drier. Pre-emergence weed control may be limited on Krasnozem or red brown Ferrosol soil types, where moisture is not maintained in the top 5 cm of soil. Refer to Soil Type in the General Instructions. Contact Sipcam for advice on variety susceptibility before using this product (refer to the CROP SAFETY section of the label). |
| Peanuts (Qld, NSW and NT only) | Awnless barnyard grass (<i>Echinochloa colona</i>), Barnyard grass (<i>E. crus-galli</i>), Blackberry nightshade (<i>Solanum nigrum</i>), Blue billygoat weed (<i>Ageratum houstonianum</i>), Common sida (<i>Sida rhombifolia</i>), Giant/Black pigweed (<i>Triathema portulacastrum</i>), Glossy nightshade (<i>Solanum americanum</i>), Green amaranth (<i>Amaranthus viridis</i>), Green summer grass (<i>Brachiaria subquadripara</i>), Ipomoea sop., Guinea grass (<i>Panicum maximum</i>), Liverseed grass (<i>Urochloa panicoides</i>), Milkweed (Mexican fire weed) (<i>Euphorbia heterophylla</i>), Pigweed (<i>Portulaca oleracea</i>), Summer grass (<i>Digitaria ciliaris</i>) | 300 to 400 mL | Pre-emergence application. Optimum weed control will occur when moisture is maintained in the top 5 cm of soil. Apply to weed-free soil prior to weed germination. Sufficient rainfall, or overhead irrigation, is required after application and prior to weed emergence to wet soil to a minimum depth of 5 cm. Use the higher rates for increased residual control of weeds or anticipated high weed pressure. Pre-emergence weed control may be limited on Krasnozem or red brown Ferrosol soil types, where moisture is not maintained in the top 5 cm of soil. Refer to Soil Type in the General Instructions. DO NOT apply to any one crop more than once per season. |
| | As above plus Nutgrass* (<i>Cyperus rotundus</i>) | 400 mL plus Hasten* or Kwicken* at 1 L/100 L (or a similar esterified oil adjuvant) | Post-emergence application. Apply prior to crop flowering. DO NOT apply to any one crop more than once per season. * Nutgrass: suppression only when applied to nutgrass at the 3 to 4 leaf stage. |

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:

HARVEST: NOT REQUIRED WHEN USED AS DIRECTED.

GRAZING/STOCKFEED: Wheat, Barley, Chickpea, Peanuts: DO NOT GRAZE TREATED VEGETATION, STUBBLE OR WEEDS, OR CUT FOR STOCK FOOD, FOR 4 WEEKS AFTER APPLICATION.

Sugarcane: DO NOT GRAZE CROP OR SPRAYED VEGETATION, OR CUT FOR STOCKFOOD, FOR 6 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

All Applications

Reduce the incidence of "fines" by using the correct nozzles and operating pressures. Replace worn or old nozzles and calibrate equipment on a regular basis.

Fallow Application

Brimstone can be applied in fallow prior to the planting of various crops as outlined in the Follow Crops Table. Brimstone is a pre-emergence herbicide, which is recommended to provide short term (6 to 12 weeks) preemergence control of summer germinating grasses and broadleaved weeds in cultivated or uncultivated fallow. A green weed cover of greater than 50% will significantly reduce the length of pre-emergence weed control.

Sugarcane Application

Pre-emergence

Where possible, apply to soil two days prior to incorporation. Incorporate with adequate irrigation or rainfall to a minimum depth of 5 cm. Incorporation by flood irrigation may result in loss of weed control. Brimstone should be applied to plant or ratoon cane before the cane has emerged. When applying Brimstone in moist conditions, allow weeds to germinate prior to application and apply mixed with an appropriate knockdown herbicide. Whenever emerged weeds are present, Brimstone should be mixed with an appropriate knockdown herbicide.

Post-emergence

Brimstone MUST always be mixed with registered rates of paraquat and applied as a carefully directed inter-row spray to weeds and at the base of the sugarcane stalks, minimising spray contact with sugarcane foliage. DO NOT add crop oil to this mixture. Other surfactant or adjuvant is not required.

Peanut Application

Pre-emergence

Apply to soil and incorporate with adequate irrigation or rainfall to a minimum depth of 5 cm.

Post-emergence

Brimstone must always be mixed with an adjuvant as directed in the DIRECTIONS FOR USE section of the label.

Soil Type

Pre-emergence weed control with Brimstone can be limited when applied onto soils with pH <5.0 and/or which contain high concentrations of iron (Fe) and/or aluminium (Al). These situations are common in soil types such as Krasnozems and red brown Ferrosols.

MIXING

Brimstone is an aqueous solution. Part fill the spray tank with water, then with agitator running, add the required amount of product, then fill the tank with water. When tank mixing this product with other recommended compatible products, first add the other product to the tank and mix thoroughly before adding this product. For post-emergence applications in peanuts, add adjuvant after the addition of Brimstone to the tank. No adjuvants or surfactants are required when Brimstone is used alone for pre-emergence weed control or with paraquat. DO NOT add crop oil when applied post-emergence with paraquat.

APPLICATION

DO NOT apply by aircraft.

Fallow application

Apply with boom equipment in not less than 50 L/ha water using flat fan nozzles. Avoid overlap and do not overspray headlands.

Sugarcane:

Ensure thorough coverage of soil and weeds with well-calibrated spray equipment. Apply in a minimum of 200 L water/ha using a nozzle configuration to generate a medium to coarse spray pattern for pre-emergence applications or medium spray pattern for post-emergence applications. Replace worn or old nozzles and calibrate equipment on a regular basis.

Peanuts:

Apply with boom equipment in not less than 50 L/ha water using flat fan nozzles. Avoid overlap and do not overspray headlands. For post-emergence applications, Brimstone should be applied a minimum of 2 hours before rainfall.

EQUIPMENT CLEAN-UP

Thoroughly flush all spray equipment with water following the use of Brimstone and before use with other products.

COMPATIBILITY

Brimstone is compatible with pendimethalin, atrazine, glyphosate, paraquat, paraquat+ diquat, metsulfuron methyl, triclopyr 600, Touchdown* and 2,4-D. It is NOT compatible with dicamba.

RESISTANT WEEDS WARNING

GROUP 2 HERBICIDE

Brimstone Herbicide is a member of the imidazolinone (IMIs) group of herbicides. The product has the inhibitors of acetolactate synthase (ALS inhibitors), acetohydroxyacid synthase (AHAS) mode of action. For weed resistance management, the product is a Group 2 herbicide. Some naturally-occurring weed biotypes resistant to this product and other Group 2 herbicides, may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population, if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group 2 herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Sipcam Australia Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant weeds.

FOLLOW CROPS

Under conditions which do not favour breakdown of this product, carry-over soil residues can affect susceptible follow crops. As environmental and agronomic factors make it impossible to eliminate all risks associated with this product, rotational crop injury is always possible. The following minimum re-cropping intervals (months after application) should be observed.

Following use in summer fallow:

| MONTHS AFTER APPLICATION | | | | |
|---|--|---|-----------------------------|-----------------|
| 3 | 4 | 10 | 24 | 36 |
| Faba beans, peanuts, mungbeans, maize varieties with CLEARFIELD* Technology ONLY: -Pacific Hycorn 621T -Pacific Hycorn 531T -Pioneer 33951R Wheat varieties with CLEARFIELD Technology only | Wheat* (except for wheat varieties with CLEARFIELD* Technology ONLY), barley*, lucerne*, chickpea | Sorghum ⁼ , maize ⁼ (except for maize varieties with CLEARFIELD* Technology ONLY), dryland cotton [^] | Irrigated cotton, sunflower | Any other crops |

* The following additional requirements apply if it is intended to plant wheat (except for varieties with CLEARFIELD Technology), barley or lucerne during the next winter season.

DO NOT apply Brimstone in fallow later than the end of December.

DO NOT apply Brimstone in areas where rainfall from spraying to sowing of cereals is expected to be below 200 mm.

⁼ The following additional requirements apply if it is intended to plant sorghum or maize (except for varieties with CLEARFIELD Technology) during the next summer season.

DO NOT apply Brimstone in areas where rainfall from spraying to sowing of sorghum or maize is expected to be below 500 mm.

[^] The following additional requirements apply if it is intended to plant dryland cotton during the next summer season.

DO NOT apply Brimstone in areas where rainfall from spraying to sowing of dryland cotton is expected to be below 550 mm.

Following use in sugarcane and peanuts:

| MONTHS AFTER APPLICATION | | | | | | |
|--|--|-------------------------|--|----------------------|-----------------------------|-----------------|
| 0 | 4 | 5 | 10 | 18 | 24 | 36 |
| Mung beans, Peanuts, soybeans, maize varieties with CLEARFIELD Technology ONLY: -Pacific Hycorn 621T -Pacific Hycorn 531T -Pioneer 33951R | Chick peas, lucerne*, barley*, wheat* (except for wheat varieties with CLEARFIELD Technology ONLY) | Lupins, pasture legumes | Maize ⁼ (except varieties with CLEARFIELD Technology ONLY), dryland cotton [^] | Sorghum ⁼ | Irrigated cotton, sunflower | Any other crops |

* The following additional requirements apply if it is intended to plant wheat (except for varieties with CLEARFIELD Technology), barley or lucerne during the next winter season.

DO NOT apply Brimstone in areas where rainfall from spraying to sowing of cereals is expected to be below 200 mm.

⁼ The following additional requirements apply if it is intended to plant sorghum or maize (except for varieties with CLEARFIELD Technology), during the next summer season.

DO NOT apply Brimstone in areas where rainfall from spraying to sowing of sorghum or maize is expected to be below 500 mm.

[^] The following additional requirements apply if it is intended to plant dryland cotton during the next summer season.

DO NOT apply Brimstone in areas where rainfall from spraying to sowing of dryland cotton is expected to be below 550 mm.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. DO NOT spray within 50 m of wetlands or waterways.

CROP SAFETY

This product may, in some circumstances, lead to transient crop yellowing and temporary slowing of growth in sugarcane but plants soon recover and yield is unaffected. This effect may be more pronounced when the product is used under conditions where soil moisture is high within 2 days following application and on light/ sandy soil.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. When opening the container and preparing the spray, wear elbow length PVC gloves. Wash hands after use. After each days use, wash gloves.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. *Phone Australia 131126.*

SAFETY DATA SHEET

For further information, refer to the Safety Data Sheet (SDS) which is available from the supplier or from our web site, www.sipcam.com.au

NOTICE TO BUYER

This product must be used in accordance with the directions for use. Where this product is not a good of a kind ordinarily acquired for personal, domestic or household use or consumption under the Australian Consumer Law, Sipcam's liability for any direct, indirect or consequential loss which you incur, including, without limitation, loss or damage to crop, loss of equipment, property damage, personal injury or death is limited, at Sipcam's option, to the replacement of the product or the supply of an equivalent product. By opening this package, you agree to be bound by these terms. If you do not agree to these terms, please return the entire unopened package intact to Sipcam or the place in which you purchased this product for a full refund.

* Not a Sipcam trademark Clearfield is a registered trademark of BASF.