



MacroSol K Acetate



DIRECTIONS FOR USE

- Product should be used during ideal growing conditions to ensure effective uptake. Avoid extreme weather conditions such as heat and frost events. Spraying in the early morning or late afternoon can assist in minimising risk of rapid drying on the leaf due to evaporation. Where local experience of successful use is not available, we recommend a small scale jar test and test application prior to wide spread use. If any issues are observed do not continue with the application.
- It is recommended to add product to the tank mix after all other products have been mixed well in the tank, including most of the water.
- Do not apply by foliar application to stone fruit.
- Do not apply by foliar application to flowers or fruit of any crop.

COMPATIBILITY

Compatible with most common fertilisers and crop protection products. It is recommended to perform a jar test before mixing. Do not use with amine based products. Do not apply this product with copper fungicides by foliar application on all crops.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.

SAFETY DATA SHEET

For further information, refer to the Safety Data Sheet (SDS) which is available from the supplier or from our web site, www.sipcam.com.au

Nitrogen (N): 30 g/L as urea, Potassium (K): 270 g/L as potassium acetate.

A high analysis potassium acetate solution, free of any chloride, specifically designed for foliar, fertigation and furrow injection applications.

CROPS	RATE	COMMENTS
BROADACRE CROPS	Foliar 4 – 6 L/ha Furrow Injection 2 – 10 L/ha	Apply with a minimum of 80 L/ha water. Ensure product is injected below the seed. Cereals apply at early tillering to early boot stage. Canola apply before bolting.
LUCERNE	5 – 15 L/ha	Apply after cutting or at crown green.
COTTON	5 – 20 L/ha	Apply every 7 to 10 days as required.
BERRIES	5 – 20 L/ha	Apply every 10 to 14 days as required.
GRAPES	Foliar 5 – 10 L/ha Fertigation 80 – 140 L/ha	Apply from fruit set through veraison.
TREE CROPS	Fertigation 40 – 140 L/ha	Every 10 to 14 days based on site requirements.
VEGETABLES AND ROW CROPS	Fertigation 20 – 100 L/ha	Every 10 to 14 days based on site requirements.

CAUTION: Do not apply as a foliar application in conjunction with copper and EC fungicides. Applications in adverse spraying conditions or in conjunctions with some crop protection products may result in reductions in efficacy or phytotoxicity which may not be acceptable to growers. For most effective results apply early morning or late afternoon.

WARNING STATEMENTS

Do not swallow. The spray from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin.

Batch No:

Date of Manufacture:

This product is NOT subject to the provisions of the Australian Code for the Transport of Dangerous Goods by Road and Rail when packed in this container.

Specialist Advice In Emergency Only
1800 033 111
All Hours - Australia Wide