



# MacroSol NPK

## 1.8:22:7.5 + 1% Zinc



### DIRECTIONS FOR USE

- Product should be used during ideal growing conditions to ensure effective uptake. Avoid extreme weather conditions such as heat and frost events. Spraying in the early morning or late afternoon can assist in minimising risk of rapid drying on the leaf due to evaporation. Where local experience of successful use is not available, we recommend a small scale jar test and test application prior to wide spread use. If any issues are observed do not continue with the application.
- It is recommended to add product to the tank mix after all other products have been mixed well in the tank, including most of the water.
- Do not apply by foliar application to stone fruit.
- Do not apply by foliar application to flowers or fruit of any crop.

### COMPATIBILITY

Compatible with most common fertilisers and crop protection products. It is recommended to perform a jar test before mixing. Do not use with amine based products. Do not apply this product with copper fungicides by foliar application on all crops.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.

### SAFETY DATA SHEET

For further information, refer to the Safety Data Sheet (SDS) which is available from the supplier or from our web site, [www.sipcam.com.au](http://www.sipcam.com.au)

**Nitrogen: 18 g/L, Phosphorus: 220 g/L, Potassium: 75 g/L, Zinc: 10 g/L**

A high analysis NPK containing multiple micronutrients specifically designed for furrow injection applications in multiple crop types.

CROPS	RATE	COMMENTS
ALL CROPS	<b>Starter fertiliser</b> 10 – 50 L/ha	For furrow injection the product must be greater than 2.5 cm to the side or below the seed. Use a minimum of 10 L of water, up to 100 L of water depending on the crop type.
	<b>Fertigation</b> 10 – 60 L/ha	Apply towards the end of irrigation shift allowing time for the product to be flushed through the entire irrigation system.
	<b>Foliar</b> 3 – 10 L/ha	Broadacre crops apply with a minimum 100 L/Ha water. Horticultural crops apply with a 200 to 2000 L/ha.

**CAUTION:** Do not apply as a foliar application in conjunction with copper and EC fungicides. Applications in adverse spraying conditions or in conjunctions with some crop protection products may result in reductions in efficacy or phytotoxicity which may not be acceptable to growers. For most effective results apply early morning or late afternoon.

### WARNING STATEMENTS

Do not swallow. The spray from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin.

Batch No:

Date of Manufacture:

This product is NOT subject to the provisions of the Australian Code for the Transport of Dangerous Goods by Road and Rail when packed in this container

Specialist Advice In Emergency Only  
**1800 033 111**  
All Hours - Australia Wide



Sipcam Pacific  
Australia Pty Ltd  
Level 1,  
191 Malop Street  
Geelong  
VICTORIA 3220

# 1000 L

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**NOTICE TO BUYER** This product must be used in accordance with the directions for use. Where this product is not a good of a kind ordinarily acquired for personal, domestic or household use or consumption under the Australian Consumer Law, Sipcam's liability for any direct, indirect or consequential loss which you incur, including, without limitation, loss or damage to crop, loss of equipment, property damage, personal injury or death is limited, at Sipcam's option, to the replacement of the product or the supply of an equivalent product. By opening this package, you agree to be bound by these terms. If you do not agree to these terms, please return the entire unopened package intact to Sipcam or the place in which you purchased this product for a full refund.