

Multisol

NUTRITIONAL PLANT SULFATE FORMULATION



% w/v

% w/v

THE PRODUCT

 Multisol is a sulfate based multi element nutrition deficiency corrector for all crops. It can be used in a preventive program or at the onset of symptoms.

DIRECTIONS FOR USE

- Product should be used during ideal growing conditions to ensure effective uptake. Avoid extreme weather conditions such as heat and frost events. Spraving in the early morning or late afternoon can assist in minimising risk of rapid drving on the leaf due to evaporation. Where local experience of successful use is not available, we recommend a small scale iar test and test application prior to wide spread use. If any issues are observed do not continue with the application.
- It is recommended to add product to the tank mix after all other products have been mixed well in the tank, including most of the water.
- Do not apply by foliar application to stone fruit.
- Do not apply by foliar application to flowers or fruit of any crop.

COMPATIBILITY

Compatible with most common fertilisers and crop protection products. It is recommended to perform a iar test before mixing. Do not use with amine based products. Do not apply **Multisol** with copper fungicides by foliar application on all crops.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre, Phone Australia 131 126.

SAFETY DATA SHEET

For further information, refer to the Safety Data Sheet (SDS) which is available from the supplier or from our web site, www.sipcam.com.au

High sulfate analysis ZMC formulation

Total copper (Cu) content of % w/v

Total zinc (Zn) content of

% w/v

Total manganese (Mn) content of

Total sulphate (S) content of

Formulation: A multi nutrient sulfate liquid designed for furrow injection, foliar and fertigation applications where a high analysis and minimal compatibility is required to prevent or correct a nutrient deficiency.

CROPS	RATE	COMMENTS
BROADACRE CROPS	Foliar 2 – 3 L/ha Furrow Injection 4 – 10 L/ha Seed Treatment 5 L/T Soil Drench 5-12 L/ha	Apply Cereals 3-5 leaf stage. Pulses/Lucerne 10-14 days before flower Canola 4-9 true leaves Furrow Injection apply 2cm below seed depth Seed treatment apply at seed cleaning Soil Drench apply pre-seeding
TREES (excluding stonefruit), NUTS, BERRIES AND VINES	Foliar 1 – 3 L/ha Fertigation 2 – 5 L/ha	Apply in a water volume of 400L to 1,000L per ha. Apply at regular intervals of 7-14 days from petal fall up until harvest.
VEGETABLES Cucurbits Fruiting Vegetables Potatoes Carrots, Onions	Foliar & Fertigation 500mL – 2 L/ha	Apply from onset of active growth
conditions or in conjunction	s with some crop protection	junction with copper and EC fungicides. Applications in adverse spraying on products may result in reductions in efficacy or phytotoxicity which may not apply early morning or late afternoon.
		n irritant. Avoid

Specialist Advice In Emergency Only This product is NOT subject to the provisions of the Australian Code for the Transport of Dangerous Goods 1800 033 111 by Road and Rail when packed in this container All Hours - Australia Wide

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

NOTICE TO BUYER This product must be used in accordance with the directions for use. Where this product is not a good of a kind ordinarily acquired for personal, domestic or household use or consumption under the Australian Consumer Law, Sipcam's liability for any direct, indirect or consequential loss which you incur, including, without limitation, loss or damage to crop, loss of equipment, property damage, personal injury or death is limited, at Sipcam's option, to the replacement of the product or the supply of an equivalent product. By opening this package, you agree to be bound by these terms. If you do not agree to these terms, please return the entire unopened package intact to Sipcam or the place in which you purchased this product for a full refund.



