

**POISON**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# **Protek®**

## **250 EC Fungicide**

**ACTIVE CONSTITUENT: 250 g/L PROTHIOCONAZOLE**  
**SOLVENT: 175 g/L N,N-DIMETHYLDECANAMIDE**

<b>GROUP</b>	<b>3</b>	<b>FUNGICIDE</b>
--------------	----------	------------------

Systemic fungicide for the control of various diseases of Wheat, Barley, Oats and Triticale as per the Directions for Use

APVMA Approval Number: 88919/130784

Contents: 5 -1000L



**SIPCAM PACIFIC AUSTRALIA PTY LIMITED**  
**Level 1, 191 Malop Street Geelong VIC 3220**

## Protek® 250 EC Fungicide

### STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product.

### Refillable Containers

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

### SAFETY DIRECTIONS

Will irritate the eyes. May irritate the skin. Avoid contact with eyes and skin. When using together with other products, consult their safety directions. When opening the container and preparing the spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves and face shield or goggles. When applying by boomspray equipment, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing). Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

### SAFETY DATA SHEET

For further information, refer to the Safety Data Sheet (SDS) which is available from the supplier or from our web site, [www.sipcam.com.au](http://www.sipcam.com.au).

### NOTICE TO BUYER

This product must be used in accordance with the directions for use. Where this product is not a good of a kind ordinarily acquired for personal, domestic or household use or consumption under the Australian Consumer Law, Sipcam's liability for any direct, indirect or consequential loss which you incur, including, without limitation, loss or damage to crop, loss of equipment, property damage, personal injury or death is limited, at Sipcam's option, to the replacement of the product or the supply of an equivalent product. By opening this package, you agree to be bound by these terms. If you **DO NOT** agree to these terms, please return the entire unopened package intact to Sipcam or the place in which you purchased this product for a full refund.

Not subject to the provisions of the Australian Code for the Transport of Dangerous Goods by Road and Rail when packed in this container and transported in Australia.

For specialist advice in emergency only, call **1800 033 111** all hours, Australia wide

### G.H.S STATEMENT

**Hazard :** H315 Causes skin irritation. H319 Causes serious eye irritation.

**Response:** IF exposed: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention. IF ON SKIN: Wash with plenty of soap and water. If skin irritation occurs: Get medical advice/attention. Take off contaminated clothing and wash before reuse

Date of Manufacture:

Batch No:



**POISON**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# **Protek<sup>®</sup>**

## **250 EC Fungicide**

**ACTIVE CONSTITUENT: 250 g/L PROTHIOCONAZOLE**  
**SOLVENT: 175 g/L N,N-DIMETHYLDECANAMIDE**

<b>GROUP</b>	<b>3</b>	<b>FUNGICIDE</b>
--------------	----------	------------------

Systemic fungicide for the control of various diseases of Wheat, Barley, Oats and Triticale as per the Directions for Use

**IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT**

APVMA Approval Number: 88919/130784



**SIPCAM PACIFIC AUSTRALIA PTY LIMITED**  
**Level 1, 191 Malop Street Geelong VIC 3220**

## RESTRAINTS

**DO NOT** apply more than 500mL/ha per crop.

**DO NOT** apply if heavy rain has been forecasted within 48 hours.

**DO NOT** apply to waterlogged soil.

**DO NOT** irrigate past the point of runoff for 48 hours after application.

**DO NOT** apply product by spraying equipment carried on the back of the user.

## SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at

[apvma.gov.au/spraydrift](http://apvma.gov.au/spraydrift).

**DO NOT** allow bystanders to come into contact with the spray cloud.

**DO NOT** apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer zone table/s below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

**DO NOT** apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

**DO NOT** apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

**DO NOT** apply by a boom sprayer unless the following requirements are met:

- Spray droplets not smaller than a MEDIUM spray droplet size category
- Minimum distances between the application site and downwind sensitive areas (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for boom sprayers') are observed.

## Buffer zones for boom sprayers

Application rate	Boom height above the target canopy	Mandatory downwind buffer zones				
		Bystander areas	Natural aquatic areas	Pollinator areas	Vegetation areas	Livestock areas
PROTEK at up to 250 mL/ha	0.5 m or lower	0 metres	0 metres	0 metres	0 metres	0 metres
	1.0 m or lower	0 metres	15 metres	0 metres	0 metres	0 metres
PROTEK at up to 250 mL/ha with LAGUNA 430 at up to 150 mL/ha	0.5 m or lower	0 metres	0 metres	0 metres	0 metres	0 metres
	1.0 m or lower	0 metres	15 metres	0 metres	0 metres	0 metres
PROTEK at up to 250 mL/ha with OVERTURE at up to 500 mL/ha	0.5 m or lower	0 metres	0 metres	0 metres	0 metres	0 metres
	1.0 m or lower	0 metres	20 metres	0 metres	0 metres	0 metres
PROTEK at up to 250 mL/ha with ACCOLADE at up to 640 mL/ha	0.5 m or lower	0 metres	0 metres	0 metres	0 metres	0 metres
	1.0 m or lower	0 metres	20 metres	0 metres	0 metres	0 metres

**DO NOT** apply by aircraft unless the following requirements are met:

- Spray droplets not smaller than a MEDIUM spray droplet size category.
- For maximum release height above the target canopy of 3 metres or 25 per cent of wingspan or 25 per cent of rotor diameter, whichever is the greatest, minimum distances between the application site and downwind sensitive areas (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for aircraft') are observed.

### Buffer zones for aircraft

Application rate	Type of aircraft	Mandatory downwind buffer zones				
		Bystander areas	Natural aquatic areas	Pollinator areas	Vegetation areas	Livestock areas
PROTEK at up to 250 mL/ha	Fixed-wing	0 metres	60 metres	0 metres	10 metres	0 metres
	Helicopter	5 metres	50 metres	0 metres	15 metres	0 metres
PROTEK at up to 250 mL/ha with LAGUNA 430 at up to 150 mL/ha	Fixed-wing	0 metres	60 metres	0 metres	10 metres	0 metres
	Helicopter	5 metres	50 metres	0 metres	15 metres	0 metres
PROTEK at up to 250 mL/ha with OVERTURE at up to 500 mL/ha	Fixed-wing	0 metres	90 metres	0 metres	10 metres	0 metres
	Helicopter	5 metres	70 metres	0 metres	15 metres	0 metres
PROTEK at up to 250 mL/ha with ACCOLADE at up to 640 mL/ha	Fixed-wing	0 metres	75 metres	0 metres	10 metres	0 metres
	Helicopter	5 metres	60 metres	0 metres	15 metres	0 metres

## DIRECTIONS FOR USE

**Table 1: Protek Alone**

Apply at the first signs of disease. May be used twice, particularly if infection occurs early. Do not apply more than 500 mL/ha Protek per crop.

Crop	Disease	Rate mL/ha	Critical Comments
Wheat	Leaf rust ( <i>Puccinia recondita</i> f.sp. <i>tritici</i> , <i>Puccinia triticina</i> )	250	Apply at the first signs of disease. May be used twice, particularly if infection occurs early.
	Septoria nodorum - glume blotch ( <i>Phaeosphaeria nodorum</i> )		Monitor crops from late tillering. Aim to protect the three top leaves of the plant from disease. Apply with a suitable adjuvant (refer to Use of Adjuvant)
	Septoria tritici blotch ( <i>Zymoseptoria tritici</i> )		Monitor crops from mid tillering. Apply at the first sign of disease development. Monitor and reapply within 14 to 21 days if conditions favour disease development. Apply with a suitable adjuvant (refer to Use of Adjuvant).
	Powdery mildew ( <i>Blumeria graminis</i> f.sp. <i>tritici</i> )		
Barley	Net form net blotch ( <i>Pyrenophora teres</i> f. <i>teres</i> )	250	Monitor crops from mid tillering. On susceptible varieties apply at the first sign of disease development. Monitor and reapply within 14 to 21 days if conditions favour disease development. Apply with a suitable adjuvant (refer to Use of Adjuvant).
	Powdery mildew ( <i>Blumeria graminis</i> f.sp. <i>hordei</i> )		Monitor crops from mid tillering.
	Leaf scald ( <i>Rhynchosporium secalis</i> )		Monitor crops from mid tillering (earlier if no effective seed treatment has been applied). On susceptible varieties apply at the first sign of disease development. Monitor and reapply within 14 to 21 days if conditions favour disease development. Apply with a suitable adjuvant (refer to Use of Adjuvant).
	Leaf rust ( <i>Puccinia hordei</i> )		Monitor crops from late tillering. Apply at the first sign of disease development. Monitor and reapply within 14 to 21 days if conditions favour disease development. Apply with a suitable adjuvant (refer to Use of Adjuvant).
Oats	Leaf rust ( <i>Puccinia coronata</i> f.sp. <i>avenae</i> )	250	Monitor crops from early tillering and on susceptible varieties apply at the first sign of infection. -Continue to monitor crops after application. Re-application may be required if conditions favour disease development. Apply with a suitable adjuvant (refer to Use of Adjuvant).
	Septoria blotch ( <i>Phaeosphaeria avenaria</i> ) Red leather leaf ( <i>Spermospora avenae</i> )		

**Table 2: Protek + Laguna 430 Fungicide (430 g/L tebuconazole)**

Observe the most conservative WHPs, re-application intervals and other instructions on the label of this and the mixture product. DO NOT use Protek + Laguna on oats.

Crop	Disease	Rate mL/ha	Critical Comments
Wheat	Stripe rust ( <i>Puccinia striiformis</i> )	125 to 250 Protek	Monitor crops from early stem elongation, and on susceptible varieties apply at the first sign of infection. Use the high rate in higher yielding crops where conditions favour disease development or susceptible varieties are grown.
	Stem rust ( <i>Puccinia graminis tritici</i> )	<b>Plus</b>	Continue to monitor crops after application, re-application may be required if conditions favour disease development and initial application is made before the flag leaf has emerged.
	Leaf rust ( <i>Puccinia recondita</i> f.sp. <i>tritici</i> , <i>Puccinia triticina</i> )		
	Fusarium head blight/head scab ( <i>Fusarium graminearum</i> )	75 to 150 Laguna	Apply as a preventative spray at the first sign of flowering. Spray equipment must be set up to achieve good coverage of wheat heads. Use the high rate in higher yielding crops where conditions favour disease development or susceptible varieties are grown.
	Yellow leaf spot ( <i>Pyrenophora tritici-repentis</i> )		Monitor crops from late tillering and spray before disease has infected any of the top three leaves of the crop. Aim to protect the three top leaves of the plant from disease.
	Septoria nodorum - glume blotch ( <i>Phaeosphaeria nodorum</i> )		Monitor crops from late tillering. Aim to protect the three top leaves of the plant from disease. Where lower rates are used apply with a suitable adjuvant (refer to Use of Adjuvant)
	Powdery mildew ( <i>Blumeria graminis</i> f.sp. <i>tritici</i> )		Monitor crops from mid tillering. Apply at the first sign of disease development. Monitor and reapply within 14 to 21 days if conditions favour disease development. Use the high rate where conditions favour severe disease, or disease is established in the lower canopy. Where lower rates are used apply with a suitable adjuvant (refer to Use of Adjuvant).
Barley	Net form net blotch ( <i>Pyrenophora teres</i> f. <i>teres</i> )	125 to 250 Protek	Monitor crops from mid tillering. On susceptible varieties apply at the first sign of disease development. Monitor and reapply within 14 to 21 days if conditions favour disease development. Use the high rate where conditions favour severe disease.
	Spot form net blotch ( <i>Pyrenophora teres</i> f. <i>maculata</i> )	<b>Plus</b>	Where lower rates are used apply with a suitable adjuvant (refer to Use of Adjuvant).
	Powdery mildew ( <i>Blumeria graminis</i> f.sp. <i>hordei</i> )	75 to 150 Laguna	Monitor crops from mid tillering. Use the higher rate in higher yielding crops where conditions favour disease development or susceptible varieties are grown.
	Leaf scald ( <i>Rhynchosporium secalis</i> )		Monitor crops from mid tillering (earlier if no effective seed treatment has been applied). On susceptible varieties apply at the first sign of disease development. Monitor and reapply within 14 to 21 days if conditions favour disease development. Use the higher rates where conditions favour severe disease. Where lower rates are used apply with a suitable adjuvant (refer to Use of Adjuvant).
	Leaf rust ( <i>Puccinia hordei</i> )		Monitor crops from late tillering. Apply at the first sign of disease development. Monitor and reapply within 14 to 21 days if conditions favour disease development. Use the high rate where conditions favour severe disease, or disease is established in the lower canopy. Where lower rates are used apply with a suitable adjuvant (refer to Use of Adjuvant).
Triticale	Stripe rust ( <i>Puccinia striiformis</i> )	125 to 250 Protek  <b>Plus</b>  75 to 150 Laguna	Monitor crops from early stem elongation, and on susceptible varieties apply at the first sign of infection. Use the higher rate in higher yielding crops where conditions favour disease development or susceptible varieties are grown. Continue to monitor crops after application. Re-application may be required if conditions favour disease development and initial application is made before the flag leaf has emerged.

**Table 3: Protek + Overture Fungicide (125 g/L epoxiconazole)**

Add Overture to Protek for improved control of certain disease compared to Protek alone. Apply at the first signs of disease. Protek at 250mL/ha plus Overture at 250mL/ha may be used twice, particularly if infection occurs early. Use Protek plus Overture at 500mL/ha just once under higher disease pressure or when longer disease control is required. Do not apply more than 500 mL/ha of Protek or 500 mL/ha of Overture per crop. Observe the most conservative WHPs, re-application intervals and other instructions on the label of this and the mixture product.

Crop	Disease	Rate mL/ha	Critical Comments
Wheat	Leaf rust ( <i>Puccinia recondita</i> f.sp. <i>tritici</i> , <i>Puccinia triticina</i> )	250 Protek  <b>Plus</b>  500 Overture	Apply when conditions favour disease development and preferably prior to development of disease in the crop. Aim to apply between stem elongation and ear emergence complete (ZGS 32-59) if disease is present. DO NOT apply later than ZGS 59.
	Septoria nodorum - glume blotch ( <i>Phaeosphaeria nodorum</i> )	250 Protek  <b>Plus</b>  250 or 500 Overture	Monitor crops from early stem elongation, and on susceptible varieties apply at the first sign of infection. Use the high rate tank mix in higher yielding crops where conditions favour disease development or susceptible varieties are grown. Continue to monitor crops after application, re-application may be required if conditions favour disease development and initial application is made before the flag leaf has emerged.
	Powdery mildew ( <i>Blumeria graminis</i> f.sp. <i>tritici</i> )	250 Protek  <b>Plus</b>  250 Overture	Apply when conditions favour disease development and prior to development of disease in the crop. Aim to apply during early tillering (ZGS 21-22) stage of the crop if disease is present. DO NOT apply later than ZGS 59. Repeat spraying may be required particularly if infection pressure persists. Regularly monitor crops from 3-4 weeks after the first application for signs of reinfection.
Barley	Leaf rust ( <i>Puccinia hordei</i> )	250 Protek  <b>Plus</b>  250 or 500 Overture	Apply when conditions favour disease development and preferably prior to development of disease in the crop. Aim to apply between stem elongation and ear emergence complete (ZGS 32-59). DO NOT apply later than ZGS 59. Apply the higher rate tank mix when disease is present on the top leaf or conditions are favourable for disease development at these stages of crop development.
	Net form net blotch ( <i>Pyrenophora teres</i> f. <i>teres</i> )		Apply when conditions favour disease development and prior to development of disease in the crop. Aim to apply between stem elongation and ear emergence complete (ZGS 32-59) if disease is present. DO NOT apply later than ZGS 59. Two applications of Overture Fungicide at the lower rate may provide better control than a single application under certain conditions. If applying a 2-spray program apply 250mL/ha per application, the first application at ZGS 31-32 if the disease is present. Aim to apply the second application around ZGS 37-39 but no later than ZGS 59, if infection pressure persists.
	Leaf scald ( <i>Rhynchosporium secalis</i> )  Powdery mildew ( <i>Blumeria graminis</i> f.sp. <i>hordei</i> )	250 Protek  <b>Plus</b>  250 Overture	Apply when conditions favour disease development and prior to development of disease in the crop. Aim to apply between early tillering and flag leaf emergence (ZGS21-39) if disease is present. DO NOT apply later than ZGS 59. Repeat spraying may be required, particularly if infection pressure persists. Regularly monitor the crop from 3-4 weeks after the first application for signs of re-infection.
Oats	Septoria avenae blotch ( <i>Parastagonospora avenaria</i> f. sp. <i>avenaria</i> )	250 Protek  <b>Plus</b>  250 or 500 Overture	Monitor crops from mid tillering and apply at the first sign of disease development. Monitor and re-apply within 14 to 21 days if conditions favour disease development. Use the high rate tank mix where conditions favour severe disease, or disease is established in the lower canopy. Where lower rates are used apply with a suitable adjuvant (refer to Use of Adjuvant).



**Table 4: Protek + Accolade 250 EC Fungicide (250 g/L azoxystrobin)**

Add Accolade to Protek for improved control of certain disease compared to Protek alone. Apply at the first signs of disease. Protek at 250mL/ha plus Accolade at 320mL/ha may be used twice, particularly if infection occurs early. Use Protek plus Accolade at 640mL/ha just once under higher disease pressure or when longer disease control is required. Do not apply more than 500 mL/ha of Protek or 640 mL/ha of Accolade per crop. Observe the most conservative WHPs, re-application intervals and other instructions on the label of this and the mixture product.

Crop	Disease	Rate mL/ha	Critical Comments
Wheat	Septoria tritici blotch ( <i>Zymoseptoria tritici</i> )	250 Protek	Monitor crops from mid tillering and apply at the first sign of disease development.
	Yellow leaf spot ( <i>Pyrenophora tritici-repentis</i> )	<b>Plus</b> 320 or 640mL Accolade	
Barley	Net form net blotch ( <i>Pyrenophora teres f. teres</i> )		Monitor and reapply within 14 to 21 days if conditions favour disease development. Use the high rate tank mix where conditions favour severe disease, or disease is established in the lower canopy. Where lower rates are used apply with a suitable adjuvant (refer to Use of Adjuvant).
	Spot form net blotch ( <i>Pyrenophora teres f. maculata</i> )		
	Leaf rust ( <i>Puccinia hordei</i> )		
Oats	Septoria avenae blotch ( <i>Parastagonospora avenaria f. sp. avenaria</i> )		

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

#### WITHHOLDING PERIODS

##### Harvest

##### **DO NOT HARVEST FOR 5 WEEKS AFTER APPLICATION**

WHEN USING PROTEK IN A TANK MIX WITH ANOTHER PRODUCT  
OBSERVE WHICHEVER PRODUCT HARVEST WITHHOLDING PERIOD IS THE LONGER.

##### Grazing

##### **DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION**

WHEN USING PROTEK IN A TANK MIX WITH ANOTHER PRODUCT  
OBSERVE WHICHEVER PRODUCT GRAZING AND STOCK FOOD WITHHOLDING PERIOD IS THE LONGER.

#### EXPORT OF TREATED PRODUCE

Growers should note that MRLs or import tolerances do not exist in all markets for produce treated with prothioconazole. If you are growing produce for export, please check with Sipcam Pacific Australia Pty Ltd for the latest information on MRLs and import tolerances before using this product.

#### GENERAL INSTRUCTIONS

Protek is a systemic fungicide for the control of various diseases of wheat, barley, oats and triticale. Monitor the crop regularly for symptoms of disease. Apply Protek at the first sign of disease, although this will depend on factors such as expected weather conditions and the crop's resistance to disease. Protek may be tank-mixed with Laguna 430 SC, Overture Fungicide or Accolade 250 EC Fungicide for improved disease control or control of a wider spectrum of diseases. Refer to the Directions for Use for specific disease recommendations and compatibility. Ensure good coverage of all susceptible plant parts.

#### Application

**Ground:** Apply product using a spray volume of 70 – 100 L/ha and spray droplets not smaller than a MEDIUM spray droplet size category.

**Aerial:** Apply product using a minimum spray volume of 20 L/ha and spray droplets not smaller than a MEDIUM spray droplet size category.

## **FUNGICIDE RESISTANCE WARNING**

### **GROUP 3 FUNGICIDE**

Protek 250 EC is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group 3 fungicide. Some naturally occurring individual fungi resistant to the product and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Sipcarn Pacific Australia Pty Ltd accepts no liability for any losses that result from failure of these products to control resistant fungi.

### **Resistance Management**

Protek should be applied in a spray program containing fungicides with different modes of action and integrated management practices to reduce the risk of fungicide resistance. Apply Protek alone or in recommended tank-mixtures as specified in the Directions for Use, in association with CropLife Fungicide Resistance Management Strategies. Refer to the website- <https://www.croplife.org.au/resources/programs/resistance-management/>

### **USE OF ADJUVANT**

The addition of an adjuvant may improve control. Add 250mL/100L of a 1000 g/L alcohol alkoxyolate non-ionic surfactant eg BS1000\*.

### **COMPATIBILITY/TANK MIXING**

Protek is compatible with Laguna 430 SC, Overture Fungicide, Accolade 250 EC Fungicide and Slipstream 250 EC Fungicide. If tank mixes are to be used observe all directions, precautions and limitation on all products to be used. As formulations of other manufacturer's products are beyond the control of Sipcarn Pacific Australia Pty Ltd, and water quality varies with location, all mixtures should be tested prior to mixing commercial quantities.

### **PRECAUTIONS**

#### **RE-ENTRY**

Do not enter treated areas until spray has dried. If prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves and footwear. Clothing must be laundered after each day's use.

### **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

Very toxic to aquatic life. DO NOT contaminate streams, rivers or watercourses with the chemical or used containers. A spray drift minimisation strategy should be employed at all times. Spray drift may occur under adverse meteorological conditions or from certain spraying equipment. Do not allow spray to drift onto sensitive areas including, but not limited to, susceptible plants/crops, cropping land, pasture, natural streams, rivers, wetlands, waterways or human dwellings. Integrated pest management – where IPM is practiced: Protek 250 EC Fungicide may have adverse effects on some non-target beneficial insects such as predatory mites.

### **STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product.

#### **Refillable Containers**

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

## **SAFETY DIRECTIONS**

Will irritate the eyes. May irritate the skin. Avoid contact with eyes and skin. When using together with other products, consult their safety directions. When opening the container and preparing the spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves and face shield or goggles. When applying by boomspray equipment, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing). Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

## **FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

## **SAFETY DATA SHEET**

For further information, refer to the Safety Data Sheet (SDS) which is available from the supplier or from our web site, [www.sipcam.com.au](http://www.sipcam.com.au).

## **NOTICE TO BUYER**

This product must be used in accordance with the directions for use. Where this product is not a good of a kind ordinarily acquired for personal, domestic or household use or consumption under the Australian Consumer Law, Sipcam's liability for any direct, indirect or consequential loss which you incur, including, without limitation, loss or damage to crop, loss of equipment, property damage, personal injury or death is limited, at Sipcam's option, to the replacement of the product or the supply of an equivalent product. By opening this package, you agree to be bound by these terms. If you **DO NOT** agree to these terms, please return the entire unopened package intact to Sipcam or the place in which you purchased this product for a full refund.