

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

ROVER® SYSTEMIC INSECTICIDE

ACTIVE CONSTITUENT: 400g/L DIMETHOATE
(An anti-cholinesterase compound)

GROUP	1B	INSECTICIDE
-------	-----------	-------------

For the control of wide range of insect pests on fruits and vegetable crops, pastures, cotton, lucerne, oil seed crops, peanuts and ornamentals as listed in the Directions for Use table

**IMPORTANT: READ THESE INSTRUCTIONS BEFORE USING OR OTHERWISE HANDLING
THE PRODUCT**

Contents: 20 Litres

APVMA Approval No: 65259/RV0317P



SIPCAM PACIFIC AUSTRALIA PTY LIMITED
Level 1, 191 Malop Street Geelong VIC 3220

® Registered Trademark of Sipcam Pacific Australia Pty Ltd

ROVER SYSTEMIC INSECTICIDE

GENERAL INSTRUCTIONS

Before opening, carefully read Directions for Use, Precautionary Statements, Safety Directions and First Aid Instructions.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well ventilated area out of direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Protect from direct sunlight and temperatures above 40°C. If storing for more than 2-3 months, avoid temperatures above 30°C. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt

SAFETY DIRECTIONS

Poisonous if absorbed by skin contact, inhaled or swallowed. Repeated minor exposure may have a cumulative poisoning effect. Will damage eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist, a washable hat, a PVC or rubber apron, elbow-length chemical resistant gloves, a face shield and impervious footwear. When using the prepared spray (or dip for pre-plant and post-harvest dipping) wear elbow-length chemical resistant gloves. If applying by hand by vehicle mounted low pressure equipment wear cotton overalls buttoned to the neck and wrist, elbow-length chemical resistant gloves and a half face-piece respirator with organic vapour/gas cartridge or canister. If clothing becomes contaminated with product remove clothing immediately. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use wash gloves, face-shield, respirator and contaminated clothing.

FIRST AID

If swallowed, splashed on skin or in eyes, or inhaled, contact a Poisons Information Centre - Phone Australia 131 126 or a doctor at once. Remove any contaminated clothing and wash skin thoroughly. If swallowed, activated charcoal may be advised. Give atropine if instructed.

SAFETY DATA SHEET

For further information refer to Safety Data Sheet (SDS) which is available from the supplier- or by visiting:- www.sipcam.com.au.

NOTICE TO BUYER

This product must be used in accordance with the directions for use. Where this product is not a good of a kind ordinarily acquired for personal, domestic or household use or consumption under the Australian Consumer Law, Sipcam's liability for any direct, indirect or consequential loss which you incur, including, without limitation, loss or damage to crop, loss of equipment, property damage, personal injury or death is limited, at Sipcam's option, to the replacement of the product or the supply of an equivalent product. By opening this package, you agree to be bound by these terms. If you do not agree to these terms, please return the entire unopened package intact to Sipcam or the place in which you purchased this product for a full refund.

Batch Number:

Date of Manufacture:

Expiry Date:

This product is Classified as a Dangerous Good under the Australian Code for the Transport of Dangerous Goods by Road and Rail.	
UN NO. 3017	ORGANOPHOSPHORUS PESTICIDE, LIQUID, TOXIC, FLAMMABLE, N.O.S.
PG III	HAZCHEM: 3W
For specialist advice in emergency only, call 1800 033 111 all hours, Australia wide	



G.H.S. STATEMENTS

HAZARDS H315: Causes skin irritation. H320: Causes eye irritation. H335: May cause respiratory irritation. H373: May cause damage to organs through prolonged or repeated exposure.

PREVENTION P260: Do not breathe vapours or spray. P270: Do not eat, drink or smoke when using this product. P271: Use only outdoors or in a well ventilated area.

RESPONSE P301+P330+P331: IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. P304+P340: IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing. P305+P351+P338: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. P332+P337+P313: If skin irritation occurs or eye irritation persists: Get medical advice

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

ROVER®

SYSTEMIC INSECTICIDE

ACTIVE CONSTITUENT: 400g/L DIMETHOATE
(An anti-cholinesterase compound)

GROUP	1B	INSECTICIDE
-------	-----------	-------------

For the control of wide range of insect pests on fruits and vegetable crops, pastures, cotton, lucerne, oil seed crops, peanuts and ornamentals as listed in the Directions for Use table

**This leaflet is part of the label.
READ THESE INSTRUCTIONS BEFORE USING OR OTHERWISE HANDLING
THE PRODUCT**

APVMA Approval No: 65259/ RV0317P



SIPCAM PACIFIC AUSTRALIA PTY LIMITED
Level 1, 191 Malop Street Geelong VIC 3220

® Registered Trademark of Sipcam Pacific Australia Pty Ltd

DIRECTIONS FOR USE:**Restraints**

DO NOT apply to any non-food tree crop or plantation (including *Eucalyptus* spp.) by air.

DO NOT apply by misting or fogging equipment

DO NOT apply with air blast spray equipment unless operators are protected by engineering controls such as enclosed cabs fitted with appropriate air filters

DO NOT use open mixing/loading systems for aerial application

CROP	PEST CONTROLLED	STATE	RATE	WHP	CRITICAL COMMENTS
Adzuki Beans, Cowpea, Mung Bean, Navy Bean, Pigeon Pea	Thrips	NSW, WA only	800mL/ha	14 days (H) 14 days (G)	Two treatments between pre-bloom (just before flower bud formation) and pod initiation may be necessary. If infestation is severe or prolonged apply the first spray at the pre-bloom stage and the second spray at full-flowering. DO NOT re-apply within 14 days. Use sufficient water to give good coverage. Toxic to bees. DO NOT spray plants in flower while bees are foraging.
Adzuki Bean	Leafhoppers, Bean Fly, Bean Blossom Thrips	Qld, WA only	800mL/ha or 75mL/100L water	14 days (H) 14 days (G)	Apply when insects appear and repeat as necessary. For Green Vegetable Bug apply at first flowering and repeat 3 weeks later. DO NOT re-apply within 14 days. Do not spray when bees are foraging.
Avocados	Queensland Fruit Fly		75mL/100 litres of water	7 days	Apply as pest populations indicate.
Avocados (Dip)		NSW, WA only	Charge the dip at the rate of 100mL/100 litres of water	nil	Dip the fruit for 1 minute and allow to drain before packing.
Kiwifruit (Chinese Gooseberries) (Inedible peel varieties only), Lychees, (Dipping)					Dip the fruit for 1 minute and allow to drain before packing.

CROP	PEST CONTROLLED	STATE	RATE	WHP	CRITICAL COMMENTS
Bananas (dipping use only)	Fruit Fly	NSW, WA only	Charge the dip at the rate of 100mL/100 litres of water	nil	Dip fruit for 10-60 seconds. Top with concentration of 125-150mL/100L.
Beans (green vegetable)	Spider Mites, Thrips, Jassids, Green Vegetable Bug, Aphids	All States	75mL/100 litres of water	7 days (G) 7 days (H)	Apply when pests appear and repeat as necessary. For Green Vegetable Bug in beans apply at first flowering and repeat 3 weeks later. QLD ONLY: Some strains of spider mites may be resistant to organophosphorous compounds.
	Bean Fly				Apply 3 and 7 days after crop emergence and then at weekly intervals until blossom.
	Redlegged Earthmite	NSW, Vic, SA, Tas, WA only			Apply when pests appear and repeat as necessary
Beans, Peas (green vegetable – not snow or sugar snap peas)	Cow Pea Aphid	NSW, WA only	350-650 mL/ha		Apply when pests appear. Use the higher rate in cold weather.
Beetroot	Leafmining Fly		800mL/ha or 75mL/100 litres of water	Beetroot 14 days	Apply when pest damage first appears. Repeat spray if necessary.
Blueberries, bilberries and other vaccinium berries	Aphids, Thrips, Jassids, Spidermites, Redlegged earth mites	All states	75mL/100 litres of water	1 day	DO NOT exceed a maximum number of 7 applications per crop per season with a minimum 21 day interval between applications. Apply in the afternoon when bees are finished foraging. Apply when pest first appears and repeat at 3 weekly intervals or as necessary. Some strains of spidermites are resistant to organophosphorous compounds.
	Strawberry Bug, Rutherglen Bug	Qld, Vic, SA, WA, Tas only			
	Queensland Fruit Fly	NSW, WA only			
Blackberries, raspberries	Aphids, Thrips, Jassids, Spidermites, Redlegged Earthmite	All States		7 days	Apply when pest first appears and repeat at 3 weekly intervals or as necessary. Some strains of spidermites are resistant to organophosphorous compounds.
	Strawberry Bug, Rutherglen Bug	Qld, Vic, SA, WA, Tas only			
Capsicums	Aphids, Thrips, Green Vegetable Bug, Jassids	All States		3 days	Apply when pests appear. DO NOT USE as a post-harvest or post-harvest quarantine treatment. Apply when pests first appear and repeat as required
	Cucumber Fly	NSW, WA only	750mL/ha or 75mL/100L water		
	Fruit Fly				

CROP	PEST CONTROLLED	STATE	RATE	WHP	CRITICAL COMMENTS
Cereals (including maize and sorghum)	Lucerne Flea	WA, Vic, Tas, SA, NSW, only	55-85 mL/ha	4 wks (H) 14 days (G)	Apply 3-5 weeks after the commencement of autumn rains or when outbreak occurs. Use the higher rate in cold weather. Do not spray on bare ground. Allow the crop to emerge before application. Boom spray-apply in 50-100 litres/ha of water. Aircraft and misting machines 20-40 litres/ha.
	Redlegged Earthmite	WA, Vic, Tas, SA only			
		NSW only	85 mL/ha		
	Wingless Grasshopper	All States	75mL/100 litres of water or 750mL/ha		Apply when grasshoppers appear and re-apply as required. In addition to the infested area spray a band of about 20 metres around areas to be protected.
	Brown Wheat Mite	Qld, WA only	90mL/ha		Apply when mites appear.
	Blue Oat Mite	Qld, NSW, WA only			
	Leafhoppers, Cereal Aphids	Qld, WA only	500mL/ha		Apply when pests threaten damage to crop.
Citrus Fruit (except Meyer Lemons, Seville Oranges & Cumquats)	Queensland Fruit Fly	NSW, Vic, Qld, WA only	75mL/100 litres of water	7 days	Qld, NSW, Vic only: Do not use on Meyer lemons, Seville oranges and cumquats. Apply 2 full cover sprays 2 weeks apart 7 weeks and 5 weeks before harvest. If harvesting is delayed a third spray may be required.
	Mediterranean Fruit Fly	NSW, Vic, WA only			
	Aphids, Thrips	All States			Apply when pests appear
	Bronze Orange Bug	NSW, Vic, Qld, SA, WA only			Apply when pests appear and as necessary
	Wingless Grasshopper	All States			Apply when grasshoppers appear and re-apply as required. In addition to the infested area spray a band of about 20 metres around the areas to be protected.
Cowpeas	Aphids, Mirid Bugs	Qld, WA only	500mL/ha	14 days (G)	Apply when pests appear. DO NOT re-apply within 14 days.
	Bean Fly, Leafhoppers		800mL/ha	14 days (H)	
	Thrips		500mL/ha		

CROP	PEST CONTROLLED	STATE	RATE	WHP	CRITICAL COMMENTS	
Cotton	Lucerne Flea	WA, NSW, SA, Tas only	55-85mL/ha	14 days	Apply 3-5 weeks after the commencement of autumn rains or when outbreak occurs. Use the higher rate in cold weather. Do not spray on bare ground. Allow the crop to emerge before application. Boom spray - apply in 50-100 litres/ha of water. Aircraft - apply in 20-40 litres/ha.	
	Redlegged Earthmite	WA, Vic, Tas, SA only				
		NSW only	85 mL/ha			
	Aphids, Spider Mites (incl Red Spider, Twospotted Mite)	Qld, NSW, WA only	500mL/ha	Apply when pests appear and repeat as required. Use the higher rate for heavy infestations. NOTE: Some strains of spider mites are resistant to organophosphorous compounds. DO NOT use this product where resistant strains are present. DO NOT feed cotton fodder, stubble or trash to livestock.		
	Thrips		350-375 mL/ha			
	Wingless Grasshopper		750mL/ha or 75mL/100L water			
	Leafhoppers (Jassids)		350mL/ha			
Bugs (inc. Green Vegetable Bug, Green Mirids, Apple Dimpling Bug, Brown Smudge Bug, Broken-backed Bug, Rutherglen Bug)	340-500mL/ha					
Melons including watermelons	Cucumber Fly	NSW, Qld, WA only	75mL/100 litres of water or 750mL/100 litres per ha Low volume		7 days	Apply when pests appear and repeat as necessary
Zucchini	Aphids, Thrips, Green Vegetable Bug, Jassids	All States			1 day	
Custard Apple (Dip)	Queensland Fruit Fly	NSW, WA only	Charge the dip at the rate of 100mL/100 litres of water	nil	Dip the fruit for 1 minute and allow to drain before packing.	
Duboisia	Thrips	Qld, WA only	75mL/100 litres water as an overall spray	nil	Apply every 7-10 days or as pest population indicates.	
Grain Legumes	Spider Mites, Thrips, Jassids, Green Vegetable Bug, Aphids, Bean Fly	Qld, Vic, SA, WA, Tas only	75mL/100L or 800mL/ha	14 days (H)	Apply when insects appear and repeat as necessary. DO NOT re-apply within 14 days. Spray when flowering spikes carrying 20-50 aphids that are easy to find, and when there is evidence of virus disease. Some strains of spider mites are resistant to organophosphorous compounds.	
	Redlegged Earthmite	Vic, SA, Tas, WA only	75mL/100L	14 days (G)		
	Lucerne Flea	WA only	85mL/100L			Apply at emergence.

CROP	PEST CONTROLLED	STATE	RATE	WHP	CRITICAL COMMENTS
Lentils	Redlegged Earth Mite	NSW, WA only	90mL/ha	14 days (H) 14 days (G)	Apply when pests first appear. DO NOT re-apply within 14 days
Litchi	Litchi Erinose Mite	Qld, NSW, WA only	75mL/100L water		Pre-planting dip: Immerse plants in mixture for 1 minute and drain before planting in the field. Established trees: Apply just before a growth flush and repeat at 14-21 day intervals until all new growth is damage-free
Lucerne	Lucerne Flea	WA, Vic, Tas, NSW, SA only	55-85 mL/ha	-	Apply 3-5 weeks after the commencement of autumn rains or when outbreak occurs. Use higher rate in cold weather or on mature pasture. Do not spray on bare ground. Allow the lucerne to emerge before application. Boom spray - apply in 50-100 litres of water/ha. Aircraft and misting machines - apply in 20-40 litres/ha
	Redlegged Earthmite	WA, Vic, Tas SA only	55-85 mL/ha		
		NSW only	85 mL/ha		
Maize	Maize Leafhopper, Thrips	Qld, WA only	500 mL/ha	4 weeks (H) 14 days (G)	Apply 2 sprays 5-7 days apart.
Mangoes	Queensland Fruit Fly	Qld, NSW, Vic, WA only	75 mL/100 litres of water	3 days,	Apply as a cover spray at first sign of infestation
	Mediterranean Fruit Fly	WA, Vic, NSW only			
Mangoes, Pawpaw, Passionfruit (Dips)	Queensland Fruit Fly	NSW, WA only	Charge the dip at the rate of 100 mL/100 litres of water	nil	Dip the fruit for 1 minute and allow to drain before packing.
Mung Beans	Aphids, Mirid Bugs	Qld, WA only	500 mL/ha	14 days (H) 14 days (G)	Apply when pests first appear. DO NOT re-apply within 14 days.
	Bean Fly, Leafhoppers		800 mL/ha		
Navy Beans	Aphids, Mirid Bugs	Qld, NSW, WA Only	500 mL/ha		

CROP	PEST CONTROLLED	STATE	RATE	WHP	CRITICAL COMMENTS
Oil Seeds (other than cotton and peanuts)	Lucerne Flea	WA, NSW, SA, Tas only	55-85 mL/ha	-	Apply 3-5 weeks after the commencement of autumn rains or when outbreak occurs. Use the higher rate in cold weather. Do not spray on bare ground. Allow the crop to emerge before application. DO NOT apply more than 7 days after crop emergence. Boom spray - apply in 50-100 litres/ha of water. Aircraft - apply in 20-40 litres/ha.
	Redlegged Earthmite	WA, Vic, Tas, SA only			
		NSW only	85 mL/ha		
Ornamentals (not Chrysanthemums, Begonias, Liquid Amber or Gloxinias)	Aphids, Thrips, Jassids, Spider Mites, Leafhoppers, Azalea Lace Bug, Green Vegetable Bug, Leaf Miners, Green House White Fly	All States	75 mL/100 litres of water	nil	Apply when pests first appear and repeat as necessary. Some strains of Spider mites are resistant to organophosphorous compounds.
	Bronze Orange Bug	NSW, Qld, SA, Vic, WA only			
	Wingless Grasshopper	All States			Apply when grasshoppers appear and re-apply as required. In addition to the infested area spray a band of about 20 metres around area to be protected.
	Woolly aphid	Vic, SA, Tas, WA only			Apply when pest first appears and repeat as necessary
Wild Flowers, Protea	Aphids, Thrips, Leafhoppers, Rutherglen Bug	WA only			Apply when pests appear. Rover will not control OP resistant mites.

CROP	PEST CONTROLLED	STATE	RATE	WHP	CRITICAL COMMENTS
Pastures, Pasture Seed & Forage Crops other than cereal and grain legume crops, inc. Clover, Medics, Lucerne, Legumes (for animal feed)	Redlegged Earth Mite, Lucerne Flea	NSW, Vic, Tas, SA, WA only	55-85 mL/ha	14 days (G)	Apply 3-5 weeks after the commencement of autumn rains or when outbreak occurs. Use the higher rate in NSW, and in cold weather and/or heavy infestations in other States. Do not spray on bare ground. Allow the crop to emerge before application. DO NOT apply more than 7 days after crop emergence. Boom spray - apply in 50-100 litres of water. Aircraft - apply in 20-40 litres/ha.
Peanuts	Lucerne Flea	WA, NSW, SA, Tas only	55-85mL/ha	14 days (G)	Apply 3-5 weeks after the commencement of autumn rains or when outbreak occurs. Use the higher rate in cold weather. Do not spray on bare ground. Allow the crop to emerge before application. Boom spray - apply in 50-100 litres/ha of water. Aircraft - apply in 20-40 litres/ha.
	Redlegged Earthmite	WA, Vic, Tas, SA only		14 days (H)	
		NSW only	85 mL/ha		
	Wingless Grasshopper	All States	75 mL/100 litres of water or 750 mL/ha		Apply when grasshoppers appear and re-apply as required. In addition to the infested area spray a band of about 20 metres around areas to be protected.
	Jassids, Green Vegetable Bug,		350 mL/ha		Apply when pests appear.
Aphids, Thrips, Peanut Mite	Qld, NSW, WA only				
Peas (green vegetable – not snow or sugar snap peas)	Spider Mites, Thrips, Jassids, Green Vegetable Bug, Aphids, Bean Fly	All States	75 mL/100 litres of water	7 days (H) 7 days (G)	Apply when pests appear and repeat as necessary. For Green Vegetable Bug apply at first flowering and repeat 3 weeks later. Do not spray when bees are foraging.
	Redlegged Earth Mite	NSW, Vic, SA, Tas, WA only			

CROP	PEST CONTROLLED	STATE	RATE	WHP	CRITICAL COMMENTS
Potatoes	Thrips, Aphids, Jassids, Green Vegetable Bug	All States	75 mL/100 litres of water	14 days (H)	Apply when pests appear.
	Redlegged Earthmite	NSW, Vic, SA, Tas, WA only			Apply when pests first appear and repeat at 3 weekly intervals as required
Onions, Beetroot, Turnip	Redlegged Earthmite	NSW, Vic, SA, Tas, WA only		Onions 7 days Beetroot, Turnip 14 days	Apply when pests first appear and repeat at 3 weekly intervals as required

CROP	PEST CONTROLLED	STATE	RATE	WHP	CRITICAL COMMENTS
Soy Beans	Green Vegetable Bug, Jassids	NSW, SA, WA, Tas, Qld only	340 mL/ha	14 days (H) 14 days (G)	Apply when pests appear
Tobacco	Lucerne Flea, Redlegged Earth Mite	NSW, WA only	80 mL/100L water	4 weeks	Apply spray to tobacco in seed bed when insects are present. Re-apply after 7 days if necessary.
Tomatoes (processing only)	Queensland Fruit Fly	Qld, NSW, Vic, WA only	750 mL/ha or 75 mL/100 L water	21 days	<p>QLD ONLY: Apply full cover sprays at 4 and 3 weeks prior to harvest.</p> <p>NSW ONLY: Apply full cover sprays at 4 and 3 weeks prior to harvest.</p> <p>VIC ONLY: Apply at 7 and 5 weeks prior to harvest</p> <p>WA ONLY: Apply about 6 weeks before fruit ripens. Re-apply at fortnightly intervals. DO NOT USE on tomatoes grown in covered or protected situations such as glasshouses, greenhouses or plastic tunnels. DO NOT USE on cherry, grape or mini tomatoes</p>
	Mediterranean Fruit Fly	WA only			
	Tomato Mite	NSW, SA, Tas, Vic, WA only	60 mL/100 L		
	Bryobia Mite	SA, Tas, Vic WA only			
	Aphids, Jassids, Green Vegetable Bug, Thrips	All States	750 mL/ha or 75 mL/100 L water		Apply when insects appear. DO NOT USE on tomatoes grown in covered or protected situations such as glasshouses, greenhouses or plastic tunnels. DO NOT USE on cherry, grape or mini tomatoes

CROP	PEST CONTROLLED	STATE	RATE	WHP	CRITICAL COMMENTS
Tomatoes, large, field grown for fresh consumption	Tomato Mite	NSW, SA, Tas, Vic, WA only	60mL/100L		Apply when pest appears and repeat as necessary. DO NOT apply after commencement of flowering. DO NOT USE on tomatoes grown in covered or protected situations such as glasshouses, greenhouses or plastic tunnels. DO NOT USE on cherry, grape or mini tomatoes.
	Bryobia Mite	SA, Vic, WA, Tas only	75mL/100L		
	Aphids, Green Vegetable Bug, Jassids, Thrips	All states			
Trees: Eucalypts, Kurrajongs, Flame Trees, Umbrella Trees	Jarrah Leaf Miner, Psyllids, Kurrajongs Leaf Miner, Leaf Blister, Sawfly, Lerp Insects, Scale Insects, Spittle Bugs, Mites.	WA only	25 mL/8 litres of water	nil	Apply in early June for control of Jarrah Leaf Miner and in early Spring for Psyllids. Apply in late January as above for Kurrajongs Leaf Miner.
Vegetables:					
Zucchini	Aphids, Thrips, Jassids, Mites, Leafhoppers, Green Vegetable Bug	All States	75 mL/100 litres of water	1 day	Apply when pests appear. Rover will not control OP resistant mites.
Capsicum, chilli	Wingless Grasshoppers		750 mL/ha or 75 mL/100 L water	3 days	Apply when grasshoppers appear and re-apply as required. In addition to the infested area spray a band of about 20 meters around areas to be protected.
Asparagus, Melons including watermelon, onions, rhubarb, sweet corn, beans and-peas (green vegetable – not snow or sugar snap peas)				7 days (H), sweet corn, beans, peas 7 days (H), 7 days (G)	
beetroot, eggplant, potatoes, sweet potatoes, turnip				14 days	
tomatoes for processing, celery				21 days	
Zucchini	Cucumber Fly		NSW, WA only	75 mL/100 litres of water	1 day
Zucchini	Cucumber Fly	NSW, WA only	75 mL/100 litres of water	1 day	Apply when pests appear and repeat when necessary.

NOT TO BE USED OR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

Citrus	DO NOT harvest for 7 days after application.
Blueberries (and other vaccinium berries including bilberries)	DO NOT harvest for 1 day after application.
Blackberries, Raspberries	DO NOT harvest for 7 days after application.
Avocado, Litchi/Lychee	DO NOT harvest for 7 days after application.
Mango	DO NOT harvest for 3 days after application.
Post Harvest Dipping (Avocados, Bananas, Cactus Fruit, Chilli, Custard Apples, Feijoas, Guavas, Kiwifruit (Chinese Gooseberries inedible peel varieties), Litchi (Lychees), Mangoes, Passionfruit, Banana Passionfruit, Pawpaws, Pomegranates, Tamarillos)	Not Required When Used as Directed (dip uses only).
Litchi (pre-planting dip)	Harvest withholding period: Not required when used as directed.
Asparagus, Melons (including watermelons), Onions, Rhubarb	DO NOT harvest for 7 days after application.
Beans, Peas (green vegetables not snow or sugar snap peas)	DO NOT harvest for 7 days after application. DO NOT graze or cut for stockfood for 7 days after application.
Beetroot, Eggplant, Potatoes, Sweet Potatoes, Turnip	DO NOT harvest for 14 days after application.
Tomatoes (for processing)	DO NOT harvest for 21 days after application.
Tomatoes, large, field grown for fresh consumption	Not required when used as directed (ie DO NOT apply after commencement of flowering)
Capsicums	DO NOT harvest for 3 days after application.
Zucchini	DO NOT harvest for 1 day after application.
Cereals, (including maize, sorghum)	DO NOT harvest for 4 weeks after application. DO NOT graze or cut for stockfood for 14 days after application.
Cotton	DO NOT harvest for 14 days after application. DO NOT Feed Cotton Fodder, Stubble or Trash To Livestock.
Pulses (grain legumes)	DO NOT harvest for 14 days after application. DO NOT graze or cut for stockfood for 14 days after application.
Lucerne, Oilseeds (other than cotton and peanuts), Pastures and forage crops	Not Required When Used As Directed (ie DO NOT apply more than 7 days after crop emergence)

GENERAL INSTRUCTIONS:

Before opening, carefully read Directions for Use, Precautionary Statements, Safety Directions and First Aid Instructions.

INSECTICIDE RESISTANCE WARNING

GROUP 1B INSECTICIDE

For insecticide resistance management Rover Insecticide is a Group 1B insecticide. Some naturally-occurring insect biotypes resistant to Rover Insecticide and other Group 1B insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Rover Insecticide or other Group 1B Insecticides are used repeatedly. The effectiveness of Rover Insecticide on resistant individuals could be significantly reduced. Since the occurrence of resistant individuals is difficult to detect prior to use, Sipcam Pacific Australia Pty Ltd accepts no liability for any losses that may result from the failure of Rover Insecticide to control resistant insects. Rover Insecticide may be subject to specific resistance strategies. For further information contact your local supplier, Sipcam Pacific Australia Pty Ltd representative or local department of agriculture agronomist.

RE-ENTRY AND RE-HANDLING

Avocado, mango trees: DO NOT allow entry into treated areas for 9 days for fruit thinning and for 2– days for hand harvesting. DO NOT allow entry into treated areas for hand pruning, irrigation, orchard maintenance, weeding, scouting, or transplanting until the spray has dried. If prior entry is required, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and gloves. Clothing must be laundered after each day's use.

Citrus trees: DO NOT allow entry into treated areas for 4–days hand harvesting. DO NOT allow entry into treated areas for hand pruning, orchard maintenance, weeding, baiting/trapping, scouting, or transplanting until the spray has dried. If prior entry is required, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and gloves. Clothing must be laundered after each day's use.

Ornamentals—cut flowers or nursery plant: DO NOT allow entry into treated areas for container moving, hand harvesting of cut flowers, hand irrigation, pinching, hand pruning, scouting, transplanting, and hand weeding until the spray has dried. If prior entry is required, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and gloves. Clothing must be laundered after each day's use.

Ornamental trees farm and forest trees: DO NOT allow entry into treated areas for 9 days for hand set irrigation. DO NOT allow entry into treated areas for 7–days for hand harvesting and for 1–day for hand pruning, shaping or scouting. DO NOT allow entry into treated areas for container moving, grading/tagging, transplanting or weeding until the spray has dried. If prior entry is required, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and gloves. Clothing must be laundered after each day's use.

Glasshouses and other confined areas: DO NOT re-enter until spray deposits have dried and areas has been thoroughly ventilated.

All other crops (litchi, blackberries, raspberries, vegetables, grain legumes, cereals, cotton, oilseeds, forage crops, tobacco, ornamental shrubs, duboisia, oil tea tree): DO NOT enter treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

Post-harvest dipping of fruit and vegetables, and pre-plant dipping of plants: DO NOT handle treated fruit, vegetable or plant until the product solution has dried. If prior handling is required, wear elbow-length chemical resistant gloves

PROTECTION OF LIVESTOCK

Dangerous to bees. DO NOT spray any plants in flower while bees are foraging.

PROTECTIONS OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well ventilated area out of direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Protect from direct sunlight and temperatures above 40°C. If storing for more than 2-3 months, avoid temperatures above 30°C. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Poisonous if absorbed by skin contact, inhaled or swallowed. Repeated minor exposure may have a cumulative poisoning effect. Will damage eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist, a washable hat, a PVC or rubber apron, elbow-length chemical resistant gloves, a face shield and impervious footwear. When using the prepared spray (or dip for pre-plant and post-harvest dipping) wear elbow-length chemical resistant gloves. If applying by hand by vehicle mounted low pressure equipment wear cotton overalls buttoned to the neck and wrist, elbow-length chemical resistant gloves and a half face-piece respirator with organic vapour/gas cartridge or canister.

If clothing becomes contaminated with product remove clothing immediately. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use wash gloves, face-shield, respirator and contaminated clothing.

FIRST AID

If swallowed, splashed on skin or in eyes, or inhaled, contact a Poisons Information Centre - Phone Australia 131 126 or a doctor at once. Remove any contaminated clothing and wash skin thoroughly. If swallowed, activated charcoal may be advised. Give atropine if instructed.

SAFETY DATA SHEET

For further information refer to Safety Data Sheet (SDS) which is available from the supplier or by visiting:- www.sipcam.com.au.

NOTICE TO BUYER

This product must be used in accordance with the directions for use. Where this product is not a good of a kind ordinarily acquired for personal, domestic or household use or consumption under the Australian Consumer Law, Sipcam's liability for any direct, indirect or consequential loss which you incur, including, without limitation, loss or damage to crop, loss of equipment, property damage, personal injury or death is limited, at Sipcam's option, to the replacement of the product or the supply of an equivalent product. By opening this package, you agree to be bound by these terms. If you do not agree to these terms, please return the entire unopened package intact to Sipcam or the place in which you purchased this product for a full refund.