

Zincsol

NUTRITIONAL PLANT SULFATE FORMULATION



THE PRODUCT

 Zincsol is a zinc nutrition deficiency corrector for all crops. It can be used in a preventive program or at the onset of symptoms.

DIRECTIONS FOR USE

- Product should be used during ideal growing conditions to ensure effective uptake. Avoid extreme weather conditions such as heat and frost events.
 Spraying in the early morning or late afternoon can assist in minimising risk of rapid drying on the leaf due to evaporation. Where local experience of successful use is not available, we recommend a small scale jar test and test application prior to wide spread use. If any issues are observed do not continue with the application.
- It is recommended to add product to the tank mix after all other products have been mixed well in the tank, including most of the water.
- Do not apply by foliar application to stone fruit.
- Do not apply by foliar application to flowers or fruit of any crop.

COMPATIBILITY

Compatible with most common fertilisers and crop protection products. It is recommended to perform a jar test before mixing. Do not use with amine based products. Do not apply **Zincsol** with copper fungicides by foliar application on all crops.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.

SAFETY DATA SHEET

For further information, refer to the Safety Data Sheet (SDS) which is available from the supplier or from our web site, **www.sipcam.com.au**

High sulfate analysis zinc formulation

Total zinc (Zn) content of 16.7% w/v
Total sulphate (S) content of 8.2% w/v

Formulation: A zinc sulfate liquid designed for furrow injection, foliar and fertigation applications where a high analysis and minimal compatibility is required to prevent or correct a nutrient deficiency.

CROPS	RATE	COMMENTS
BROADACRE CROPS	Foliar 2 – 3 L/ha Furrow Injection 4 – 8 L/ha Seed Treatment 5 L/T Soil Drench 5-12 L/ha	Apply Cereals 3-5 leaf stage. Pulses/Lucerne 10-14 days before flower Canola 4-9 true leaves Furrow Injection apply 2cm below seed depth Seed treatment apply at seed cleaning Soil Drench apply pre-seeding
TREES (excluding stonefruit), NUTS, BERRIES AND VINES	Foliar 1 – 3 L/ha Fertigation 2 – 5 L/ha	Apply in a water volume of 400L to 1,000L per ha. Apply at regular intervals of 7-14 days from petal fall up until harvest.
VEGETABLES Cucurbits Fruiting Vegetables Potatoes Carrots, Onions	Foliar & Fertigation 500mL – 2 L/ha	Apply from onset of active growth

CAUTION: Do not apply as a foliar application in conjunction with copper and EC fungicides. Applications in adverse spraying conditions or in conjunctions with some crop protection products may result in reductions in efficacy or phytotoxicity which may not be acceptable to growers. For most effective results apply early morning or late afternoon.

WARNING STATEMENTS

Do not swallow. The spray from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin.

Date of Manufacture:

This product is NOT subject to the provisions of the Australian Code for the Transport of Dangerous Goods by Road and Rail when packed in this container

Specialist Advice In Emergency Only 1800 033 111

All Hours - Australia Wide



Sipcam Pacific
Australia Pty Ltd
Level 1,
191 Malop Street
Geelong

1000 L